Innate Immune System

- The innate (natural) immune system is thought to constitute an **evolutionarily older** defense strategy.

- Innate immune system provides **immediate defense** against infection, and is found in all classes of plant and animal life.

- It is the **dominant immune system** found in plants, fungi, insects, and in primitive multicellular organisms.

- A defining aspect of the innate immune system is that it carries **no memory** of an encounter with a foreign organism or molecule.

Major functions of innate immune system

1. **Recruiting immune cells** to sites of **infection** and **inflammation** through the production of **cytokines**.

Inflammation is a protective response involving immune cells, blood vessels, and molecular mediators.

Cytokines are any of a number of substances which are secreted by certain cells of the immune system and have an effect on other cells.

2. **Activation of the complement cascade** to identify bacteria, activate cells and to promote clearance of dead cells or antibody complexes.

Complement cascade, is a part of the immune system that enhances (complements) the ability of antibodies and immune cells to clear microbes and damaged cells.

3. **The identification and removal of foreign substances** present in organs, tissues, the blood and lymph, by specialized white blood cells (**phagocytes** or eating cells).

4. **Activation of the adaptive immune system** through a process known as **antigen presentation** carried out by **antigen presenting cells (APCs)**.

TABLE 1-2 Summary of nonspecific host defenses

Type	Mechanism
Anatomic barriers	
Skin	Mechanical barrier retards entry of microbes. Acidic environment (pH 3–5) retards growth of microbes.
Mucous membranes	Normal flora compete with microbes for attachment sites and nutrients. Mucus entraps foreign microorganisms. Cilia propel microorganisms out of body.
Physiologic barriers	
Temperature	Normal body temperature inhibits growth of some pathogens. Fever response inhibits growth of some pathogens.
Low pH	Acidity of stomach contents kills most ingested microorganisms.
Chemical mediators	Lysozyme cleaves bacterial cell wall. Interferon induces antiviral state in uninfected cells. Complement lyses microorganisms or facilitates phagocytosis. Toll-like receptors recognize microbial molecules, signal cell to secrete immunostimulatory cytokines. Collectins disrupt cell wall of pathogen.
Phagocytic/endocytic barriers	Various cells internalize (endocytose) and break down foreign macromolecules. Specialized cells (blood monocytes, neutrophils, tissue macrophages) internalize (phagocytose), kill, and digest whole microorganisms.
Inflammatory barriers	Tissue damage and infection induce leakage of vascular fluid, containing serum proteins with antibacterial activity, and influx of phagocytic cells into the affected area.

Inflammation

Many infections, especially from small wounds are eliminated by the **combination of complement and recruitment of phagocytes**.

Five Characteristic signs of inflammation:

1. **Redness**
2. **Swelling**
3. **Heat**
4. **Pain**
5. **Loss of function**

Causative events:

- -- Vasodilation, increased capillary permeability
- -- Emigration of phagocytes
- -- Chemical mediators (chemokines)

Cells of the innate immune response

Cells of the innate system recognize, and respond to pathogens in a **generic** (general), **non-specific way**.

- All **white blood cells (WBC)** or **leukocytes** are major cells of **innate** and **adaptive** immune response.

- Leukocytes are able to **move freely** and interact and capture cellular debris, foreign particles, or invading microorganisms.

    Cellular debris: Organic waste left over after a cell dies

- Most innate immune leukocytes **cannot divide or reproduce** on their own.



Tennis ball like, round cells are white blood cells. Disc shaped cells are red blood cells.

The innate non-phagocytic leukocytes include:

Natural killer cells, mast cells, eosinophils, and **basophils.**

The innate phagocytic cells include:

Macrophages, neutrophils and **dendritic cells**.

These cells function within the immune system by **identifying** and **eliminating** pathogens that might cause infection.



Lymphocytes



A moving leukocyte with an extended hand to capture foreign materials, e.g., bacteria

Innate non-phagocytic leukocytes

Natural killer cells

- **Natural killer cells**, or **NK cells** are lymphocytes that **attack host cells** that have been infected by microbes, but **do not directly** attack invading microbes.

  For example, NK cells attack and destroy tumor cells, and virus-infected cells, through a process known as **"missing-self."**

  This term **"missing-self"** describes cells with **low levels** of a cell-surface marker called **MHC I**.



NK cell attacking an infected cell

Mast cells:

- Mast cells are a type of innate immune cell that resides in the **connective tissue** and in the **mucous membranes**.
- They are intimately associated with defense against pathogens, wound healing, but are also often associated with **allergy** and **anaphylaxis**.

Connective tissue that connects, supports, binds, or separates other tissues or organs.

Allergy is a damaging immune response by body to a substance to which it has become hypersensitive.

Anaphylaxis is an acute allergic reaction which may lead to death.

- When activated, mast cells rapidly release **characteristic granules**, rich in **histamine, heparin** and several other chemicals.

- Histamine dilates blood vessels, causing the **characteristic signs of inflammation**, and recruits **neutrophils** and **macrophages**.

- Heparin is an **anticoagulant** (stops blood from clotting).

Basophils :

- These cells are among **granulocytes.**

- **Basophils** releasing **histamine** are important in defense against **parasites**, and play a role in **allergic reactions (such as asthma)**.

- They resemble **mast** cells.



Eosinophils:

- These cells are among **granulocytes.**
- Upon activation, eosinophils secrete a range of **highly toxic proteins** and **free radicals** that are highly effective in **killing bacteria and parasites**.
- They are also responsible for **tissue damage** occurring during **allergic reactions**.



Phagocytes

- The word 'phagocyte' literally means **'eating cell'.**
- To engulf a particle or pathogen, a phagocyte extends portions of its plasma membrane, **wrapping the membrane around the particle** until it is enveloped.
- Once inside the cell, the invading pathogen is contained inside an **endosome** which merges with a **lysosome** which contains **enzymes and acids** that **kill and digest** the particle or organism.
- Both **endosomes** and **lysosomes** are vacuoles.

Phagocytes generally **patrol the body** searching for pathogens, but are also able to react to **cytokines**.

A phagocyte attacking bacteria

The phagocytic cells of the immune system include:

1. **Macrophages**
2. **Neutrophils**
3. **Dendritic cells**.

Macrophages

- Macrophages meaning **"large eating cell",** are large phagocytic leukocytes.

- They are able to move outside of the vascular system by **moving across the cell membrane** of capillary vessels and entering the areas between cells in pursuit of invading pathogens.

- Macrophages are the **most efficient phagocytes**, and can phagocytose a large number of bacteria or other microbes and cells.

- The binding of bacterial molecules to receptors on the surface of a macrophage triggers it to engulf and destroy the bacteria through the release of **ROS (reactive oxygen species).**

- Pathogens also stimulate the macrophage to produce **chemokines**, which summons other cells to the site of infection.

ROS are unstable molecules that contain oxygen and that easily react with other molecules in a cell. A build up of reactive oxygen species in cells may cause damage to DNA, RNA, and proteins, and may cause cell death.

Chemokines are a family of small cytokines with ability to induce directed chemotaxis.


rbc
bacteria (green)
macrophage
http://www.transferpoint.net/assets/images/macrophage.jpg


1
Bacterium becomes attached to membrane evaginations called pseudopodia
2
Bacterium is ingested, forming phagosome
3
Phagosome fuses with lysosome
4
Lysosomal enzymes digest captured material
5
Digestion products are released from cell

Neutrophils

- Neutrophils are also among **granulocytes** due to the presence of granules in their cytoplasm.

- They are also known as **polymorphonuclear cells (PMNs)** due to their distinctive lobed nuclei which are usually more than two.



- Neutrophil granules contain a variety of **toxic substances** that kill or inhibit growth of bacteria and fungi.

- Neutrophils attack pathogens by releasing **ROS** and **hypochlorite**.

- Neutrophils are the **most abundant** type of phagocytes, normally representing **50 to 60%** of the total circulating leukocytes.

- They are usually the **first cells** to arrive at the site of an infection.



A neutrophil attacking anthrax bacilli

Dendritic cells

- Dendritic cells (DC) are phagocytic cells **present in tissues** that are in contact with the external environment, mainly the **skin** (where they are often called **Langerhans cells**), and the inner mucosal lining of the **nose, lungs, stomach** and **intestines**.

- Dendritic cells are very important in the process of **antigen presentation**, and serve as a **link between the innate** and **adaptive immune systems.**

A dendritic cell

Northern lights   (aurora borealis)


''ہم پرندوں کی طرح ہوا میں اڑنا
اور مچھلی کی طرح پانی میں
تیرنا تو سیکھ گئے ہیں، بس اب
یہ سیکھنا باقی ہے کہ زمین پر
انسانوں کی طرح
کیسے رہیں۔''
جاج برنارڈ شا


Thank You!